

# Failure of Donor Lymphocyte Infusion to Prevent Graft Rejection in Dogs Given DLA-Identical Marrow after 1 Gy of Total Body Irradiation

Frédéric Baron,<sup>1,4</sup> Brenda M. Sandmaier,<sup>1,2</sup> Eustacia Zellmer,<sup>1</sup> Mohamed Sorror,<sup>1</sup> Barry Storer,<sup>1,3</sup> Rainer Storb<sup>1,2</sup>

<sup>1</sup>Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington; <sup>2</sup>Department of Medicine, University of Washington School of Medicine, Seattle, Washington; <sup>3</sup>Department of Biostatistics, University of Washington School of Medicine, Seattle, Washington; <sup>4</sup>Department of Hematology, University of Liège, Liège, Belgium

Correspondence and reprint requests: Rainer Storb, MD, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N, D1-100, PO Box 19024, Seattle, WA 98109-1024 (e-mail: [rstorb@fhcrc.org](mailto:rstorb@fhcrc.org)).

Received February 22, 2006; accepted May 23, 2006

## ABSTRACT

We investigated in a preclinical canine model of hematopoietic cell transplantation (HCT) whether preemptive donor lymphocyte infusion (DLI) given 1 month after HCT could prevent late graft rejection that was the rule in historical dogs given suboptimal conditioning with 1 Gy of total body irradiation (TBI) before and immunosuppression with cyclosporine (CSP) and either mycophenolate mofetil (MMF;  $n = 6$ ) or rapamycin ( $n = 5$ ) after dog leukocyte antigen (DLA)-identical marrow transplantation. Nine dogs given DLA-identical marrow after 1 Gy of TBI followed by postgrafting MMF and CSP were studied. A single DLI was given 28-36 days after HCT, either with ( $n = 5$ ) or without ( $n = 4$ ) preceding treatment with the immunosuppressive drug pentostatin. Two of the 4 dogs given DLI only maintained stable mixed donor–host chimera beyond 30 weeks after HCT, whereas 2 rejected their grafts, on weeks 10 and 15 after HCT. One of the 5 dogs given pentostatin before DLI maintained a stable mixed donor–host chimera beyond 30 weeks, whereas 4 rejected their grafts, at weeks 8, 12, 12, and 16 after HCT. The 30-week probability of stable mixed chimerism was 33% among dogs given DLI, versus 0% among 11 historical dogs ( $P = .003$ ). In conclusion, DLI was only moderately effective in preventing graft rejection in this model. Additional immunosuppression with pentostatin did not improve that outcome. The model might be useful in developing potential strategies aimed at preventing graft rejection in patients with low donor chimerism levels.

© 2006 American Society for Blood and Marrow Transplantation

## KEY WORDS

Hematopoietic cell transplantation • Donor lymphocyte infusion • Pentostatin • Chimerism  
• Nonmyeloablative • Dog

## INTRODUCTION

Nonfatal late rejections of hematopoietic grafts may occur after nonmyeloablative or reduced-intensity conditioning regimens, particularly in patients with chronic myeloid leukemia or myelodysplastic syndromes given grafts from unrelated donors [1-4]. Low levels ( $\leq 50\%$ ) of donor chimerism in T cells [5,6] and natural killer (NK) cells [1,6,7] on days 14 and 28 after transplantation are predictive of a greater risk of graft rejection. Specifically, patients with donor NK- and T-cell chimerism levels  $\leq 50\%$  on day 14

after hematopoietic cell transplantation (HCT) had a 50% probability of graft rejection [6]. Assessing the efficacy of strategies aimed at preventing graft rejection in patients with low-donor chimerism levels has proven difficult, because only approximately half of the patients with low donor chimerism levels eventually experience graft rejection; this emphasizes the need for a good animal model.

We have previously shown that 92% and 86% of dogs given dog leukocyte antigen (DLA)-identical marrow after 2 Gy of total body irradiation (TBI) and postgrafting immunosuppression with cyclosporine

(CSP) plus mycophenolate mofetil (MMF) [8], or with CSP plus rapamycin [9], achieved sustained mixed donor–host chimerism. This approach has been successfully translated to treat human patients with hematologic malignancies or primary immunodeficiency diseases [2,5,10], demonstrating the clinical relevance of the canine model. However, when the TBI dose was reduced to 1 Gy, graft rejection was observed in 6 of 6 dogs given CSP and MMF [8] and in 5 of 5 dogs given CSP and rapamycin [9]. Graft rejections observed after 1 Gy of TBI in this model were consistent with host-versus-graft reactions rather than with a lack of marrow space [11–13], as reviewed by Baron and Storb [10].

Here we investigated whether a preemptive donor lymphocyte infusion (DLI) given 1 month after HCT could prevent late graft rejection in dogs given 1 Gy of TBI before and MMF and CSP after DLA-identical marrow transplantation.

## MATERIALS AND METHODS

Nine donor/recipient pairs (8 pairs of littermates and 1 pair of siblings), 7–10 (median 9) months old, and raised at the Fred Hutchinson Cancer Research Center, were DLA-identical on the basis of matching for highly polymorphic DLA-associated class I and class II microsatellite marker polymorphisms and DLA-DRB1 sequencing [9,14,15].

Recipients were administered 1 Gy of TBI delivered at 7 cGy/min from a linear accelerator (CLINAC 4; Varian, Palo Alto, CA). Marrow cells were aspirated from donors under general anesthesia through long needles inserted into the humeri. Marrow grafts containing  $2.4 \times 10^8$ – $6.3 \times 10^8$  (median,  $3.8 \times 10^8$ ) total nucleated cells per kg of the recipient were given through intravenous (IV) infusion within hours of TBI. Immunosuppression consisted of MMF, 10 mg/kg twice a day injected subcutaneously, from days 0–27, and CSP, 15 mg/kg twice a day orally, starting on day –1. CSP was discontinued either 1 day before DLI (for dogs not given pentostatin) or 1 day before pentostatin. Donor peripheral blood mononuclear cells (PBMCs) were collected 28–38 days (median, 33 days) after HCT by a COBE Spectra 6-cell separator (Grambo, Lakewood, CO) and were given through IV infusion (DLI) within hours of collection. Five dogs received posttransplantation pentostatin, 4 mg/m<sup>2</sup>/day IV for either 1 day (2 days before DLI; n = 3) or 3 days (days –4 to –2 before DLI; n = 2) (Table 1). Pentostatin was given as an IV bolus injection after infusion of 500 mL of saline solution, as reported previously [16]. Pretransplantation pentostatin (either  $3 \times 4$  mg/m<sup>2</sup> or  $6 \times 4$  mg/m<sup>2</sup> IV) administration was previously shown to prolong DLA-identical marrow engraftment in dogs given 1 Gy of TBI and posttransplantation MMF plus CSP [16]. The number of T cells in the DLIs was

**Table 1.** Duration of Mixed Donor–Host Chimerism in Dogs Given Marrow Grafts from DLA-Identical Donors after 1 Gy TBI Followed (Study Group) or Not Followed (Historical Group) by Preemptive DLI Given 1 Month after Transplantation

Dog No.	Nucleated Marrow Cells ( $\times 10^8$ /kg)	Pentostatin?	DLI: T Cells $\times 10^8$ /kg (day after HCT)	Percent Donor Chimerism in PBMC before DLI	Duration of Mixed Chimerism, Among PBMC, in Weeks (final % of donor chimerism in PBMC)
<b>Historical group</b>					
E165	4.0	No	No	/	12 (0)
E166	4.0	No	No	/	10 (0)
E202	4.1	No	No	/	3 (0)
E204	4.0	No	No	/	3 (0)
E227	4.0	No	No	/	10 (0)
E228	4.0	No	No	/	10 (0)
G092†	3.6	No	No	/	9 (0)
G111†	6.3	No	No	/	11 (0)
G151†	3.7	No	No	/	3 (0)
G156†	3.8	No	No	/	9 (0)
G167†	4.2	No	No	/	9 (0)
<b>Study group</b>					
G335	6.3	No	1.0 (36)	40.0	>32 (46.5)
G376	2.4	No	2.0 (28)	8.8	15 (0)
G329	2.4	No	2.3 (39)	7.7	10 (0)
G444	5.4	No	2.6 (29)	29.3	>35 (5.0)
G468	4.0	4 mg/m <sup>2</sup>	1.6 (38)	23.1	16 (0)
G509	3.8	4 mg/m <sup>2</sup>	1.6 (30)	38.6	12 (0)
G541	3.5	4 mg/m <sup>2</sup>	1.1 (30)	16.2	>33 (15.8)
G565	2.8	3 $\times$ 4 mg/m <sup>2</sup>	1.7 (33)	9.9	7* (0)
G582	5.4	3 $\times$ 4 mg/m <sup>2</sup>	1.4 (33)	14.8	12 (0)

\*Eight weeks in the granulocyte fraction.

†These dogs were given CSP plus rapamycin instead of CSP plus MMF as postgrafting immunosuppression.

assessed by flow cytometry using a monoclonal antibody directed against canine CD3 (CA17.6F9, IgG<sub>2b</sub>), as reported previously [15]; levels ranged from  $1.0 \times 10^8$  to  $2.6 \times 10^8$  T cells/kg (Table 1).

The presence of donor cells in PBMCs and granulocytes after HCT was assessed by fluorescent variable number tandem repeat polymerase chain reaction assays using an ABI Prism 310 Genetic Analyzer and Gene Scan 3.1 software (Applied BioSystems, Foster City, CA) [15]. The endpoint of the study was stable mixed donor–host hematopoietic chimerism beyond 30 weeks after HCT.

Data from dogs in this study were compared with those from a historical group of 11 dogs given DLA-identical marrow grafts after 1 Gy TBI and receiving postgrafting immunosuppression with either CSP and MMF or CSP and rapamycin [8,9].

The study was approved by the Institutional Animal Care and Use Committee at the Fred Hutchinson Cancer Research Center.

## RESULTS


### Engraftment, DLI, and Graft Rejection

Initial donor engraftment was documented in all dogs. Two of the 4 dogs given DLI only maintained stable mixed donor–host chimeras beyond 30 weeks after HCT, whereas 2 rejected their grafts, on weeks 10 and 15 (Table 1). We then investigated in 5 dogs whether host immunosuppression with pentostatin before DLI could allay the pending rejection by increasing the efficacy of DLI. One of the 5 dogs given pentostatin before DLI maintained a stable mixed donor–host chimera beyond 30 weeks, whereas the other 4 (including the 2 dogs given 3 doses of pentostatin) rejected their grafts, on weeks 8, 12, 12, and 16 after HCT (Table 1). Donor PBMC chimerism levels before DLI were  $28.5\% \pm 11.9\%$  in the 3 dogs with sustained engraftment, compared with  $17.2\% \pm 11.9\%$  in the 6 dogs with graft rejection ( $P = \text{not significant [NS]}$ ). The absolute lymphocyte counts before DLI were  $1203 \pm 741$  in the 3 dogs with sustained engraftment, versus  $1990 \pm 712$  in the 6 dogs with graft rejection ( $P = \text{NS}$ ).

### Toxicity

Hematologic toxicities were generally mild (Figure 1). Median platelet, neutrophil, and lymphocyte nadirs were 32,000, 2077, and 396 cells/ $\mu\text{L}$ , respectively. Changes in lymphocyte counts were not statistically different between the dogs given and not given pentostatin (data not shown).

The main nonhematologic toxicity was gastrointestinal; effects included anorexia, diarrhea, and moderate weight loss, most likely due to MMF. In addition, 1 animal (G565) developed intussusception at



**Figure 1.** (A) Median peripheral blood changes in 9 dogs given 1 Gy of TBI before and MMF/CSP plus DLI (with or without preceding pentostatin) after marrow transplantation from DLA-identical donors; (B) PBMC chimerism in 4 dogs given 1 Gy of TBI before and MMF/CSP plus DLI after marrow transplantation from DLA-identical donors; (C) PBMC chimerism in 5 dogs given 1 Gy of TBI before and MMF/CSP, pentostatin plus DLI after marrow transplantation from DLA-identical donors.

day 26 after HCT that was successfully removed surgically. Intussusception is associated with the use of CSP in dogs.

Pentostatin and DLI were well tolerated. Graft-

versus-host disease (GVHD) was not observed in any dog.

### Comparison of Graft Rejection with Historical Dogs

The numbers of nucleated marrow cells transplanted in present ( $2.4 \times 10^8/\text{kg}$ – $6.3 \times 10^8/\text{kg}$  [median,  $3.8 \times 10^8/\text{kg}$ ]) and historical dogs ( $3.6 \times 10^8/\text{kg}$ – $6.3 \times 10^8/\text{kg}$  [median,  $4.0 \times 10^8/\text{kg}$ ]) were comparable ( $P = .3$ ). There was a statistical trend for decreased risk of rejection in dogs receiving higher numbers of nucleated marrow cells (analyzed as a continuous variable;  $P = .07$ ). The 30-week probability of sustained donor engraftment was 33% in dogs given DLI, versus 0% in historical dogs ( $P = .003$  in univariate analysis and  $P = .002$  after adjusting for the number of nucleated marrow cells transplanted). After excluding the 5 historical dogs given CSP plus rapamycin as postgrafting immunosuppression, the  $P$  values were .03 and .02, respectively.

### DISCUSSION

In contrast to cancer patients undergoing nonmyeloablative HCT, who most often received various amount of cytotoxic therapy before nonmyeloablative conditioning, dogs included in the current study did not receive any chemotherapy before conditioning to HCT, making them an ideal model for assessing ways to further improve donor engraftment. In a previous study, we observed stable mixed donor–host chimerism in 5 of 8 dogs given 1 Gy of TBI followed by DLA-identical marrow plus granulocyte-colony-stimulating factor–mobilized PBMC grafts (containing  $1.9 \pm 0.5 \times 10^8$  T cells/kg), and in 1 of 7 dogs given 1 Gy of TBI followed by marrow plus CD3-depleted G-PBMCs [13], suggesting that the addition of donor T cells to the marrow grafts promoted sustained engraftment. Here we investigated whether infusion of additional donor T cells as late as 5 weeks after HCT, after discontinuation of immunosuppression, would promote stable engraftment in dogs given marrow grafts after 1 Gy of TBI.

Given in this manner, DLIs prevented graft rejection in only 2 of 4 dogs, suggesting that host T cells in the other 2 dogs were already sensitized against donor minor histocompatibility antigens when immunosuppression was discontinued and were capable of destroying both the marrow graft and the infused donor lymphocytes. The lack of uniform efficacy of DLI in the current canine model is consistent with observations made in human patients, where DLIs were often ineffective in preventing graft rejection in settings of low donor chimerism levels [17,18].

We next evaluated whether pentostatin, a potent immunosuppressive drug in both humans and dogs

[16,19], could reduce the number of sensitized host lymphocytes sufficiently to set the stage for DLI to promote stable donor engraftment. This strategy met with little success, because only 1 of 5 dogs achieved stable mixed chimerism. Perhaps pentostatin affected not only sensitized host T cells, but also regulatory T cells [20], which have been implicated in the maintenance of stable mixed chimerism after nonmyeloablative conditioning [21,22].

Pentostatin and DLI appeared to be safe, and no acute GVHD was seen in dogs that achieved sustained engraftment. This was most likely related to the mixed donor–host T-cell chimerism with relatively low donor contributions, a condition associated with a low incidence of GVHD [6].

In conclusion, DLI was only moderately effective in allaying graft rejection in this model, even when preceded by host immunosuppression with pentostatin. The canine model might be useful in developing better strategies for preventing graft rejection in patients with low donor chimerism levels. These strategies should be aimed at specifically targeting activated (sensitized) T cells, without affecting regulatory T cells. Potential candidates include bortezomib, recently shown to specifically inhibit activated T cells [23], and radionuclide-conjugated monoclonal antibodies targeting the CD70 antigen [24].

### ACKNOWLEDGMENTS

This work was supported in part by National Institutes of Health grants CA78902, CA15704, HL36444, and DK56465. Additional support was provided by grants from the Laura Landro Salomon Endowment Fund and the Jose Carreras International Leukemia Foundation. F.B. is a research associate of the National Fund for Scientific Research Belgium and is supported in part by postdoctoral grants from the Fulbright Commission and the Centre Anticancéreux près l'Université de Liège. We thank Michael A. Harkey, Clonal Analysis Core, for some of the variable number tandem repeat studies; Michele Spector, Alix Joslyn, and the technicians in the canine facilities of the Fred Hutchinson Cancer Research Center; Patrice Stroup, for the DLA typing; the technicians of the hematology and pathology laboratories of the Seattle Cancer Care Alliance; George E. Sale, for performing the pathology; Drs. Beard, Burroughs, Diaconescu, Graves, Gerull, Jochum, Kerbaux, Kiem, Mielcarek, Nash, and Wang, who participated in the weekend treatments; Murad Yunusov, for the correction of the intussusception on G565; and Bonnie Larson, Helen Crawford, and Sue Carbonneau, for help with manuscript preparation. The authors are grateful to Supergen, Montville, NJ, for the gift of pentostatin; Dr. Elizabeth Squires, SangStat Medical Corporation,



Fremont, CA, for the gift of cyclosporine; and Dr. Sabine Hadulco, Roche Bioscience, Nutley, NJ, for the gift of mycophenolate mofetil.

## REFERENCES

- Bornhauser M, Thiede C, Platzbecker U, et al. Dose-reduced conditioning and allogeneic hematopoietic stem cell transplantation from unrelated donors in 42 patients. *Clin Cancer Res*. 2001;7:2254-2262.
- Maris MB, Niederwieser D, Sandmaier BM, et al. HLA-matched unrelated donor hematopoietic cell transplantation after nonmyeloablative conditioning for patients with hematologic malignancies. *Blood*. 2003;102:2021-2030.
- Baron F, Maris MB, Storer BE, et al. HLA-matched unrelated donor hematopoietic cell transplantation after nonmyeloablative conditioning for patients with chronic myeloid leukemia. *Biol Blood Marrow Transplant*. 2005;11:272-279.
- Hallemeier C. Re: High rate of graft failure in 25 patients with chronic myelogenous leukemia conditioned with a reduced-intensity regimen of 550-cGy total body irradiation and cyclophosphamide for unrelated donor transplantation [letter]. *Biol Blood Marrow Transplant*. 2004;10:726-727.
- McSweeney PA, Niederwieser D, Shizuru JA, et al. Hematopoietic cell transplantation in older patients with hematologic malignancies: replacing high-dose cytotoxic therapy with graft-versus-tumor effects. *Blood*. 2001;97:3390-3400.
- Baron F, Baker JE, Storb R, et al. Kinetics of engraftment in patients with hematologic malignancies given allogeneic hematopoietic cell transplantation after nonmyeloablative conditioning. *Blood*. 2004;104:2254-2262.
- Matthes-Martin S, Lion T, Haas OA, et al. Lineage-specific chimaerism after stem cell transplantation in children following reduced intensity conditioning: potential predictive value of NK cell chimaerism for late graft rejection. *Leukemia*. 2003;17:1934-1942.
- Storb R, Yu C, Wagner JL, et al. Stable mixed hematopoietic chimerism in DLA-identical littermate dogs given sublethal total body irradiation before and pharmacological immunosuppression after marrow transplantation. *Blood*. 1997;89:3048-3054.
- Hogan WJ, Little MT, Zellmer E, et al. Postgrafting immunosuppression with sirolimus and cyclosporine facilitates stable mixed hematopoietic chimerism in dogs given sublethal total body irradiation before marrow transplantation from DLA-identical littermates. *Biol Blood Marrow Transplant*. 2003 Aug;9(8):489-495.
- Baron F, Storb R. Allogeneic hematopoietic cell transplantation following nonmyeloablative conditioning as treatment for hematologic malignancies and inherited blood disorders [review]. *Mol Ther*. 2006;13:26-41.
- Storb R, Yu C, Barnett T, et al. Stable mixed hematopoietic chimerism in dog leukocyte antigen-identical littermate dogs given lymph node irradiation before and pharmacologic immunosuppression after marrow transplantation. *Blood*. 1999;94:1131-1136.
- Storb R, Yu C, Zaucha JM, et al. Stable mixed hematopoietic chimerism in dogs given donor antigen, CTLA4Ig, and 100-cGy total body irradiation before and pharmacologic immunosuppression after marrow transplant. *Blood*. 1999;94:2523-2529.
- Zaucha JM, Zellmer E, Georges G, et al. G-CSF-mobilized peripheral blood mononuclear cells added to marrow facilitates engraftment in nonmyeloablative canine recipients: CD3 cells are required. *Biol Blood Marrow Transplant*. 2001;7:613-619.
- Diaconescu R, Little M-T, Leisenring W, et al. What role is there for antithymocyte globulin in allogeneic nonmyeloablative canine hematopoietic cell transplantation? *Biol Blood Marrow Transplant*. 2005;11:335-344.
- Burroughs L, Mielcarek M, Little M-T, et al. Durable engraftment of AMD3100-mobilized autologous and allogeneic peripheral blood mononuclear cells in a canine transplantation model. *Blood*. 2005;106:4002-4008.
- Panse JP, Storb R, Storer B, et al. Prolonged allogeneic marrow engraftment following nonmyeloablative conditioning using 100-cGy total body irradiation and pentostatin before and pharmacological immunosuppression after transplantation. *Transplantation*. 2005;80:1518-1521.
- Dey BR, McAfee S, Colby C, et al. Impact of prophylactic donor leukocyte infusions on mixed chimerism, graft-versus-host disease, and antitumor response in patients with advanced hematologic malignancies treated with nonmyeloablative conditioning and allogeneic bone marrow transplantation. *Biol Blood Marrow Transplant*. 2003;9:320-329.
- Bethge WA, Hegenbart U, Stuart MJ, et al. Adoptive immunotherapy with donor lymphocyte infusions after allogeneic hematopoietic cell transplantation following nonmyeloablative conditioning. *Blood*. 2004;103:790-795.
- Bolanos-Meade J, Jacobsohn DA, Margolis J, et al. Pentostatin in steroid-refractory acute graft-versus-host disease. *J Clin Oncol*. 2005;23:2661-2668.
- Bagasra O, Howedy A, Pomerantz RJ. Adenosine-deaminase-associated immunodeficiency. I. Differential sensitivities of lymphocyte subpopulations exposed to 2-deoxycytosine in vivo. *Clin Exp Immunol*. 1992;88:383-388.
- Georges GE, Storb R, Thompson JD, et al. Adoptive immunotherapy in canine mixed chimeras after nonmyeloablative hematopoietic cell transplantation. *Blood*. 2000;95:3262-3269.
- Lesnikova M, Nikitine A, Pogosov L, et al. Peripheral CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cells (Treg) block alloreactive host anti-donor T cells in canine mixed hematopoietic chimeras [abstract]. *Blood*. 2005;106(Pt 1):867a.
- Blanco B, Pérez-Simón JA, Sánchez-Abarca LI, et al. Bortezomib induces selective depletion of alloreactive T lymphocytes and decreases the production of Th1 cytokines. *Blood*. 2006;107:3575-3583.
- Yamada A, Salama AD, Sho M, et al. CD70 signaling is critical for CD28-independent CD8<sup>+</sup> T cell-mediated alloimmune responses in vivo. *J Immunol*. 2005;174:1357-1364.